

CASE REPORT

Open Access

Protein-C deficiency presenting as pulmonary embolism and myocardial infarction in the same patient

Syed Maqbool^{1*}, Vishal Rastogi¹, Ashok Seth², Satbir Singh³, Vijay Kumar¹ and Arif Mustaqueem⁴

Abstract

Plasma protein-C exerts anticoagulatory effects by inactivating factors V and VIII. Hereditary protein C deficiency is transmitted as an autosomal dominant disorder. Homozygous individuals usually develop purpura fulminans as newborns; heterozygous protein C-deficient individuals are at increased risk for venous thrombosis and pulmonary embolism. However, arterial thrombosis occurring as a result of congenital protein-C deficiency is still controversial. We describe a young patient with heterozygous protein-C deficiency who experienced both pulmonary embolism as well as myocardial infarction due to thrombotic occlusion without underlying major risk factors. Acute myocardial infarction in young without underlying major risk factors may be evaluated for protein c deficiency.

Introduction

Protein C is a 62-kD, vitamin K-dependent glycoprotein synthesized in the liver. It circulates in the blood as an inactive zymogen at a concentration of 4 µg/ml. Its activation into the serine-protease enzyme activated protein C (aPC), is catalyzed by thrombin when it is bound to the endothelial glycoprotein thrombomodulin [1,2]. aPC exerts its anticoagulant activity primarily through inactivation of coagulation factors Va and VIIIa, which are required for factor X activation and thrombin generation. The catalytic activity of aPC is greatly enhanced by the vitamin K-dependent cofactor protein S [3]. A deficiency of protein C disturbs the delicate balance between pro-coagulant and anticoagulant proteins and engenders a prothrombotic state. Cardinal manifestation of protein-C deficiency is venous thromboembolism. However, there are several case reports of arterial stroke [4,5] and myocardial infarction [6-11] occurring in young adults with congenital protein-C deficiency; but the results of larger studies are conflicting [12-16] and the existence of an association between protein-C deficiency and arterial thrombosis remains controversial. We present a case who had pulmonary thromboembolism 3-years back and now presented with acute myocardial infarction

without any underlying major risk factors for coronary artery disease.

Case presentation

Our patient is a 37-years old male who had presented in 2009 with acute severe dyspnea and retrosternal chest pain; his Electrocardiogram showed sinus tachycardia with s1 q3 t3 pattern; Echo showed mildly dilated right ventricle; D-dimer was >15micrograms/ml; CT angiography showed acute pulmonary thromboembolism (Figure 1) and right lower limb external iliac, common femoral and popliteal vein deep venous thrombosis (Figure 2). He was managed with heparin and discharged on oral anticoagulation. Though, erratic with his anticoagulation, he nevertheless was doing well till morning of 26th Jan 2013 when he developed severe retrosternal chest pain radiating to both arms. He was brought to hospital in an hour and his Electrocardiogram showed extensive anterior wall myocardial infarction (Figure 3). He was directly taken to the lab where the angiogram showed distal left anterior descending and proximal diagonal-2 cut off with thrombus (Figure 4). Thrombosuction was done and TIMI- 3 flow was restored (Figure 5). The underlying arteries were clean with no evidence of atherosclerosis. His electrocardiogram post thrombosuction reverted almost to normal (Figure 6). He had no major risk factors for coronary artery disease (Table 1). Patient was

* Correspondence: syedmaqbool20@yahoo.com

¹Invasive Cardiology, Fortis Escorts Heart Institute, New Delhi 110025, India
Full list of author information is available at the end of the article



Figure 5 RAO cranial view post thrombosuction showing fully patent LAD and D2 without any underlying atherosclerosis.

deficiency only six had angiography done, 2 underwent coronary intervention, one received a stent and in other the obstruction could not be relieved. Rest of the patients were thrombolysed but coronaries remained totally occluded in 3 and partially in one [6]. In our patient just thrombosuction cleaned up the coronaries and post thrombosuction even mild plaquing was not seen anywhere in the coronary tree. Our patient had absolutely no major risk factors for coronary artery disease. Patent foramen ovale was ruled out by contrast TEE. These findings suggest that protein c deficiency may cause thrombus formation in the arterial system




Figure 6 Electrocardiogram after thrombosuction.

Table 1 CAD risk factors

Blood sugar	
Fasting	86
PP	110
HbA1c	5.1%
LDL	96 mg/dl
TG	110 mg/dl
BMI	20
B.P.	120/80 mmHg
Non-smoker	

without any underlying atherosclerosis and point to the fact that young patients with acute myocardial infarction without underlying major risk factors for coronary artery diseases may be evaluated for protein c deficiency. Protein-C deficiency can occur during acute thrombosis or when the patient is on anticoagulation and the only way to confirm protein-C deficiency in such cases is to check the family members or to stop anticoagulation for two weeks and to wait until acute stage is over. Our patient had recurrent thrombotic episodes so we preferred to check the family members rather than stopping the anticoagulation. His father showed a very low level of 26% and mother had a level of 46% of protein-C activity, but there was no family history of thrombosis. It is known that all patients with protein-C and/or S deficiency donot experience thrombotic events and that 0.3% of blood donors have low protein-C levels without any overt thromboembolic episodes [21].

Table 2 Coagulation profile

Test	Patient value	Reference range
Protein-C activity, plasma	42%	67–195%
Protein-S activity, plasma	99%	77–143%
Antithrombin activity, plasma	94%	70–122%
Homocystiene, serum/plasma	11.25 micromoles	5.46 -16.20 μmoles
Factor V leiden mutation	Not Detected	
Antiphospholipid antibodies		
<i>Lupus Anticoagulant</i>		
DRWT	36 sec	32.8–48.9 sec
DRWT T: C Ratio	0.89	0.82–1.22
<i>Anticardiolipin Antibody (Aeskulia-Cardiolipin-GM) Elisa</i>		
Anticardiolipin (IgG)	1.14 GPL	POSITIVE >15.0 GPL
Anti Beta 2 Glycoprotein 1 (IgG)	1.10 GPL	POSITIVE >15.0 GPL

Conclusion

Protein-C deficiency can cause myocardial infarction in young adults with normal coronaries and without any underlying major risk factors for coronary artery disease. The message is that young patients with acute myocardial infarction with no major risk factors may be evaluated for protein c deficiency; however, it needs larger studies to be established.

Informed consent

“Written informed consent was obtained from the patient for the publication of this report and any accompanying images”.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SM conception data accumulation. AS & VR analysis. SS & AM echocardiography. VK editing. All authors have read and approved the final version of the manuscript.

Author details

¹Invasive Cardiology, Fortis Escorts Heart Institute, New Delhi 110025, India. ²Cardiovascular Sciences, Fortis Escorts Heart Institute, New Delhi 110025, India. ³Fortis Escorts Heart Institute, New Delhi 110025, India. ⁴Noninvasive Cardiology, Fortis Escorts Heart Institute, New Delhi 110025, India.

Received: 14 March 2013 Accepted: 9 September 2013

Published: 1 October 2013

References

- Dahlback B: Advances in understanding pathogenic mechanisms of thrombophilic disorders. *Blood* 2008, **112**(1):19–27.
- Clouse LH, Comp PC: The regulation of hemostasis: the protein C system. *N Engl J Med* 1986, **314**(20):1298–1304.
- Walker FJ: Regulation of activated protein C by a new protein. A possible function for bovine protein S. *J Biol Chem* 1980, **255**(12):5521–5524.
- Kohler J, Kasper J, Witt I, von Reutern GM: Ischemic stroke due to protein C deficiency. *Stroke* 1990, **21**(7):1077–1080.
- Camerlingo M, Finazzi G, Casto L, et al: Inherited protein C deficiency and non-hemorrhagic arterial stroke in young adults. *Neurology* 1991, **41**(9):1371–1373.
- Peterman MA, Roberts WC: Syndrome of protein C deficiency and anterior wall acute myocardial infarction at a young age from a single coronary occlusion with otherwise normal coronary arteries. *Am J Cardiol* 2003, **92**(6):768–770.
- Coller BS, Owen J, Jetsy J, Horowitz D, Reitman MJ, Spear J, Yeh T, Comp PC: Deficiency of plasma protein S, protein C, or antithrombin III and arterial thrombosis. *Arteriosclerosis* 1987, **7**:456–462.
- Hacker SM, Williamson BD, Lisco S, Kure J, Shoa M, Pitt B: Protein deficiency and acute myocardial infarction in the third decade. *Am J Cardiol* 1991, **68**:137–138.
- Kario K, Matsuo T, Tai S, Sakamoto S, Yamada T, Miki T, Matsuo M: Congenital protein C deficiency and myocardial infarction. *Thromb Res* 1992, **67**:95–103.
- Bux-Gewehr I, Nacke A, Feurle GE: Recurring myocardial infarction in a 35 year-old woman. *Heart* 1999, **81**:316–317.
- Sadiq A, Ahmed S, Karim A, Spivak J, Mattana J: Acute myocardial infarction: a rare complication of protein C deficiency. *Am J Med* 2001, **110**:414.
- Douay X, Lucas C, Caron C, Goudemand J, Leys D: Antithrombin, protein C and protein S levels in 127 consecutive young adults with ischemic stroke. *Acta Neurol Scand* 1998, **98**(2):124–127.
- Munts AG, van Genderen PJ, Dippel DW, van Kooten F, Koudstaal PJ: Coagulation disorders in young adults with acute cerebral ischaemia. *J Neurol* 1998, **245**(1):21–25.
- Boekholdt SM, Kramer MH: Arterial thrombosis and the role of thrombophilia. *Semin Thromb Hemost* 2007, **33**(6):588–596.
- Folsom AR, Ohira T, Yamagishi K, Cushman M: Low protein C and incidence of ischemic stroke and coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) Study. *J Thromb Haemost* 2009, **7**(11):1774–1778.
- Kenet G, Lüttkhoff LK, Albigetti M, Bernard T, Bonduel M, Brandao L, et al: Impact of thrombophilia on risk of arterial ischemic stroke or cerebral sinovenous thrombosis in neonates and children: a systematic review and meta-analysis of observational studies. *Circulation* 2010, **121**(16):1838–1847.
- Tait RC, Walker ID, Reitsma PH, et al: Prevalence of protein C deficiency in the healthy population. *Thromb Haemost* 1995, **73**(1):87–93.
- Miletich J, Sherman L, Broze G Jr: Absence of thrombosis in subjects with heterozygous protein C deficiency. *N Engl J Med* 1987, **317**(16):991–996.
- Martinelli I, Mannucci PM, De Stefano V, et al: Different risks of thrombosis in four coagulation defects associated with inherited thrombophilia: a study of 150 families. *Blood* 1998, **92**(7):2353–2358.
- Koster T, Rosendaal FR, Briet E, et al: Protein C deficiency in a controlled series of unselected outpatients: an infrequent but clear risk factor for venous thrombosis (Leiden Thrombophilia Study). *Blood* 1995, **85**(10):2756–2761.
- Gouault-Heilmann M, Leroy-Matheron C, Levent M: Inherited protein S deficiency: clinical manifestations and laboratory findings in 63 patients. *Thromb Res* 1994, **76**:269–279.

doi:10.1186/1477-9560-11-19

Cite this article as: Maqbool et al.: Protein-C deficiency presenting as pulmonary embolism and myocardial infarction in the same patient. *Thrombosis Journal* 2013 **11**:19.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

