

Original clinical investigation

Open Access

Comparison of platelet function between sedentary individuals and competitive athletes at rest

Giuseppe Lippi*¹, Martina Montagnana¹, Gian Luca Salvagno¹,
Massimo Franchini² and Gian Cesare Guidi¹

Address: ¹Sezione di Chimica e Microscopia Clinica, Università degli Studi di Verona, Verona, Italy and ²Servizio di Immunoematologia e Trasfusione, Azienda Ospedaliera di Verona, Verona, Italy

Email: Giuseppe Lippi* - ulippi@tin.it; Martina Montagnana - tina23177@interfree.it; Gian Luca Salvagno - gian_lu77@hotmail.com; Massimo Franchini - mfranchini@univr.it; Gian Cesare Guidi - gguidi@univr.it

* Corresponding author

Published: 17 August 2006

Received: 08 June 2006

Thrombosis Journal 2006, **4**:10 doi:10.1186/1477-9560-4-10

Accepted: 17 August 2006

This article is available from: <http://www.thrombosisjournal.com/content/4/1/10>

© 2006 Lippi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: There are controversial evidences on the effect of different types and workloads of physical exercise on primary hemostasis. In particular, little is known on the chronic influence of a strenuous and regular aerobic training regimen on platelet function.

Methods: The aim of this investigation was to compare platelet function between sedentary controls and trained athletes at rest and to evaluate whether a greater amount of exercise performed in professional cyclists may contribute to increased platelet chronic responsiveness compared to both elite cyclists and sedentary individuals. Platelet's ability to adhere and aggregate was assayed following a 12–24 h resting period in 49 active professional male road cyclists, 40 elite male cyclists and 43 matched sedentary healthy male volunteers, by the platelet function analyzer 100 (PFA-100).

Results and discussion: Mean values of the collagen-epinephrine test did not differ between controls and athletes (sedentary controls: 111 ± 33 s; elite athletes: 113 ± 26 s, $p = 0.93$; professional athletes: 120 ± 33 s; $p = 0.33$), whereas mean values of the collagen-ADP test displayed a slightly but significant trend towards decreased values when comparing sedentary controls (83 ± 21 s) with either elite (77 ± 11 s, $p < 0.01$) or professional (75 ± 16 s, $p < 0.01$) athletes.

Conclusion: The trend towards slightly lower collagen-ADP values are suggestive for a modest but significant chronic activation of primary hemostasis, highlighting the need to set appropriate reference ranges for the PFA-100 when evaluating primary hemostasis in physically active subjects.

Background

Recent guidelines recommend 30 minutes or more of daily, moderate-intensity physical exercise on the basis of favorable biological adaptations and substantial health benefits, which include a reduced risk of developing cardiovascular disorders, cancer and other chronic patholo-

gies [1]. Although it appears to exist a linear relation between physical activity and global health status, such that a further increase in physical activity and fitness will lead to additional improvements in health status [2], there is debate regarding intensity and type of physical activity required to achieve beneficial health changes

without overwhelming favorable outcomes and eliciting cardiovascular abnormalities not present at rest [1,3]. In particular, there are still controversial evidences on the relation between physical exercise and primary hemostasis [3-10]. Therefore, the aim of this investigation was to compare platelet function between sedentary controls and trained athletes at rest and to evaluate whether a greater amount of exercise performed in professional cyclists may contribute to influence the chronic platelet responsiveness compared to both elite cyclists and sedentary individuals.

Methods

Platelet's ability to adhere and aggregate was monitored with the use of the platelet function analyzer 100 (PFA-100, Dade International Inc. Miami, FL, USA) PFA-100 in 49 active professional male road cyclists, 40 elite male cyclists and 43 sedentary healthy male volunteers, matched for ethnic origin (all Caucasian), age (sedentary controls: 29 ± 9 y, elite cyclists: 28 ± 12 y, $p = 0.501$; professional cyclists: 28 ± 4 y, $p = 0.311$) and smoking habit (all non-smokers). Unlike platelet aggregation studies, the PFA-100 is a test relatively easy and rapid to perform, which provides a rapid evaluation of primary hemostasis and platelet function on whole blood anticoagulated specimens. The analyzer uses a disposable cartridge which internal active membrane is coated with either collagen-ADP (CADP) or collagen-epinephrine (CEPI). The interaction of whole anticoagulated blood with the agonists, in condition of high shear rates, triggers platelet activation and the resulting formation of a stable platelet plug interrupts the blood flow, which is recorded in term of closure time (CT) [11]. Athletes reached our Center in the middle of the competitive season, during a regular, high-workload period of aerobic training. All subjects recruited to the study were in a fasted state and athletes had rested for a period of 12–24 h since the last training session or competition. The consumption of drugs known to interfere with platelet function, along with clinically symptomatic platelet disorders, were carefully excluded by clinical history and physical examination in both the athlete and control populations. The research was carried out according to the principles of the Declaration of Helsinki; the study protocol was reviewed and approved by the ethics committee of our Department, and informed consent for testing was received from all individuals included in this study. Blood was collected in the morning from fasting subjects by venipuncture with 20 gauge (G) straight needles (Terumo Europe NV, Leuven, Belgium), directly into siliconized vacuum tubes containing 0.105 mol/l buffered trisodium citrate (Becton-Dickinson, Oxford, UK). Samples for PFA-100 testing were immediately analyzed, whereas specimens for coagulation studies were centrifuged at 1500 g for 10 min at 15 °C, plasma was separated, stored in aliquots and kept frozen at -70 °C until measure-

ment. Von Willebrand factor (VWF) was measured on the Mini Vidas Immunoanalyzer (bioMerieux, Marcy l'Etoile, France), by a rapid and quantitative enzyme-linked immunosorbent assay. Significance of differences between samples was assessed by Student's t-test; the level of statistical significance was set at $p < 0.05$.

Results and discussion

The average intensity of daily physical exercise was substantially different among the three study populations (sedentary controls <10 min/day, elite cyclists 30–90 min/day and pro cyclists >90 min/day). When comparing results of the PFA-100 between sedentary individuals (CADP: 83 ± 21 s; CEPI: 111 ± 33 s) and athletes, mean values of CEPI appeared substantially unvaried by the physical activity (elite athletes: 113 ± 26 s; $p = 0.93$ and professional athletes: 120 ± 33 s; $p = 0.33$), whereas mean CADP values displayed a slightly but significant trend towards decreased values in either elite (77 ± 11 s; $p < 0.01$) or professional (75 ± 16 s; $p < 0.01$) athletes (Figure 1). In athletes, and particularly in professionals, the distribution is more homogeneous for CAPD values than CEPI ones, regardless of a similar number of outliers for both CEPI and CAPD, which are correspondent to the same athletes. Results of VWF measurement did not differ significantly between controls (106 ± 14 IU/dl) and either elite (107 ± 13 IU/dl, $p = 0.258$) or professional cyclists (108 ± 14 IU/dl, $p = 0.144$).

Growing evidences indicate that benefit of physical activity on cardiovascular disease may result to some extent from effects on hemostasis [10-13]. Accordingly, for rehabilitation training it is recommended that the intensity of exercise should be distinctly below the individual anaerobic threshold. It is hence debated whether a strenuous aerobic exercise might counterbalance these advantageous modifications [1-3] and no definitive data are still available on the influence of physical exercise on primary hemostasis. Although it has been reported that acute bout of moderate-intensity exercise may reduce platelet activation [4], earlier studies observed acute platelet activation, aggregation and adhesion during physical exercise, as evaluated by ADP and collagen-induced aggregation [5], platelet granularity [6], filtagometry [7], enhanced release of soluble P-selectin and intracellular Ca^{2+} [8], downregulation of glycoprotein Ib [9] and moderate increase of platelet-leukocyte conjugates [10].

However, no data are available so far for the PFA-100 analyzer on blood samples collected from professional athletes at rest, undergoing a strenuous and regular aerobic training regimen. Although this is only a preliminary observation and even though the values of the PFA-100 observed in competitive athletes are still within normal ranges [14], the modestly decreased CADP values provide




Figure 1

Value distribution of collagen-ADP (CADP) or collagen-epinephrine (CEPI) induced aggregation on the PFA-100 system in sedentary healthy controls, elite and professional road cyclists. The central horizontal line indicates the mean value.

evidence of a slight, persistent platelet activation in elite and professional athletes, providing further insight to the controversy whether an increased physical activity may chronically influence hemostasis. In this regard, we conclude that gradually increased aerobic training regimens may produce a modest but significant chronic activation of primary hemostasis.

At variance with earlier studies, the innovative value of this investigation is represented by the original study design, involving a large subset of elite and professional endurance athletes who were tested following a 12–24 h resting period. Unfortunately, there is no definitive explanation to justify the discrepant behavior between CADP and CEPI in athletes. The activation of platelet adhesion/aggregation can be induced by means of heavy exercise, though warm-up exercise diminishes this effect in sedentary men [15]. Therefore, the continuous exercise, even of low intensity and possibly not reported by the athletes, may have contributed to the increased platelet responsiveness to CADP. Then, due to the inverse correlation between hematocrit and closure time under all circumstances, the hematocrit value is an additional variable that should be taken into consideration for the correct interpretation of PFA-100 measurements [16]. Accordingly, the different hematologic profile of competitive athletes [17] may provide a further explanation for the discrepancy observed between CADP and CEPI values in the athletes population.

Conclusion

The influence of common variables influencing PFA-100 results such as acquired or congenital platelet disorders and VWF levels has been accurately excluded in our study populations. Therefore, the trend towards slightly lower CADP values in athletes is suggestive for a moderate chronic platelet activation and deserves further scrutiny, highlighting that appropriate reference ranges for the PFA-100 may be implemented when evaluating primary hemostasis in physically active subjects.

Authors' contributions

GL: conceived of the study, participated in its design and coordination and drafted the manuscript; MM: participated in the design of the study, performed the statistical analysis and helped to draft the manuscript; GLS: recruited the study population, collected the samples and performed the laboratory testing; MF: recruited the study population and participated in the design of the study, GCG: participated in the design and coordination of the study.

References

1. Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr: **Relative intensity of physical activity and risk of coronary heart disease.** *Circulation* 2003, **107**:1110-6.

2. Warburton DER, Nicol CW, Bredin SSD: **Health benefits of physical activity: the evidence.** *CMAJ* 2006, **174**:801-9.
3. Lee KW, Lip GY: **Acute versus habitual exercise, thrombogenesis and exercise intensity.** *Thromb Haemost* 2004, **91**:416-9.
4. Wang JS, Liao CH: **Moderate-intensity exercise suppresses platelet activation and polymorphonuclear leukocyte interaction with surface-adherent platelets under shear flow in men.** *Thromb Haemost* 2004, **91**:587-94.
5. Dimitriadou C, Dessypris A, Louizou C, Mandalaki T: **Marathon run II: Effects on platelet aggregation.** *Thromb Haemost* 1977, **37**:451-5.
6. Kratz A, Wood MJ, Siegel AJ, Hiers JR, Van Cott EM: **Effects of marathon running on platelet activation markers: direct evidence for in vivo platelet activation.** *Am J Clin Pathol* 2006, **125**:296-300.
7. Li N, Wallen NH, Hjemdahl P: **Evidence for prothrombotic effects of exercise and limited protection by aspirin.** *Circulation* 1999, **100**:1374-9.
8. Wang JS, Chow SE, Chen JK: **Strenuous, acute exercise affects reciprocal modulation of platelet and polymorphonuclear leukocyte activities under shear flow in men.** *J Thromb Haemost* 2003, **1**:2031-7.
9. Kestin AS, Ellis PA, Barnard MR, Errichetti A, Rosner BA, Michelson AD: **Effect of strenuous exercise on platelet activation state and reactivity.** *Circulation* 1993, **88**:1502-11.
10. Hilberg T, Schmidt V, Glaser D, Schammne D, Losche W, Gabriel HH: **Platelet activity, sensitivity to agonist, and platelet – leukocyte conjugate formation after long-term exercise.** *Platelets* 2002, **13**:273-7.
11. Lippi G, Franchini M: **Laboratory screening for abnormalities of primary hemostasis: what's next?** *Clin Chem* 2001, **47**:2071.
12. Lippi G, Schena F, Salvagno GL, Montagnana M, Guidi GC: **D-dimer concentration is not influenced by previous physical activity.** *Thromb Res* 2006, **117**:731-2.
13. Lippi G, Salvagno GL, Montagnana M, Guidi GC: **Chronic influence of vigorous aerobic training on hemostasis.** *Blood Coagul Fibrinolysis* 2005, **16**:533-4.
14. Keresztes PA, Tazbir J: **The PFA-100: analysis and interpretation of a platelet function measurement.** *J Cardiovasc Nurs* 2005, **20**:405-7.
15. Wang JS, Yen HL, Yang CM: **Warm-up exercise suppresses platelet-eosinophil/neutrophil aggregation and platelet-promoted release of eosinophil/neutrophil oxidant products enhanced by severe exercise in men.** *Thromb Haemost* 2006, **95**:490-8.
16. Eugster M, Reinhart WH: **The influence of the haematocrit on primary haemostasis in vitro.** *Thromb Haemost* 2005, **94**:1213-8.
17. Banfi G, Mauri C, Morelli B, Di Gaetano N, Malgeri U, Melegati G: **Reticulocyte count, mean reticulocyte volume, immature reticulocyte fraction, and mean sphered cell volume in elite athletes: reference values and comparison with the general population.** *Clin Chem Lab Med* 2006, **44**:616-22.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

